

Adult—waist circumference risk indicator

Exported from METEOR (AIHW's Metadata Online Registry)

© Australian Institute of Health and Welfare 2024

This product, excluding the AIHW logo, Commonwealth Coat of Arms and any material owned by a third party or protected by a trademark, has been released under a Creative Commons BY 4.0 (CC BY 4.0) licence. Excluded material owned by third parties may include, for example, design and layout, images obtained under licence from third parties and signatures. We have made all reasonable efforts to identify and label material owned by third parties.

You may distribute, remix and build on this website's material but must attribute the AIHW as the copyright holder, in line with our attribution policy. The full terms and conditions of this licence are available at <https://creativecommons.org/licenses/by/4.0/>.

Enquiries relating to copyright should be addressed to info@aihw.gov.au.

Enquiries or comments on the METEOR metadata or download should be directed to the METEOR team at meteor@aihw.gov.au.

Adult—waist circumference risk indicator

Identifying and definitional attributes

Metadata item type:	Data Element Concept
METEOR identifier:	269670
Registration status:	Health! , Standard 01/03/2005
Definition:	An indicator of the sex specific category of risk of metabolic complications associated with excess abdominal adiposity in an adult.
Context:	Public health and health care:

This metadata item is used as an indicator of risk of metabolic complications associated with overweight and obesity including dyslipidaemia, glucose intolerance and hypertension. On a population basis there is a strong association between abdominal obesity and health risk. Body fat distribution has emerged as an important predictor of obesity-related morbidity and mortality. Abdominal obesity, which is more common in men than women, has, in epidemiological studies, been closely associated with conditions such as coronary heart disease, stroke, non-insulin dependent diabetes mellitus and high blood pressure. Waist circumference as an indicator of risk can be used:

- to indicate the prevalence of abdominal obesity and its sociodemographic distribution (problem identification);
- to evaluate health promotion and disease prevention programs (assessment of interventions);
- to monitor progress towards national public health policy;
- to ascertain determinants and consequences of abdominal obesity; and
- in nutrition and physical activity surveillance and long-term planning.

Waist circumference is a convenient and simple measurement that is unrelated to height, correlates closely with body mass index (BMI) and waist-to-hip ratio (WHR) and is an approximate index of intra-abdominal fat mass and total body fat. Changes in waist circumference can reflect changes in risk factors for cardiovascular disease and other forms of chronic disease, even though the risks seem to vary in different populations.

Object Class attributes

Identifying and definitional attributes

Object class:	Adult
Definition:	An individual aged 18 years or older.
Specialisation of:	Person/group of persons

Source and reference attributes

Submitting organisation:	Australia Institute of Health and Welfare
---------------------------------	---

Property attributes

Identifying and definitional attributes

Property:	Waist circumference risk indicator
Definition:	An indicator of the level of the risk of metabolic complications associated with excess abdominal adiposity.
Property group:	Physical wellbeing

Source and reference attributes

Submitting organisation: Australian Institute of Health and Welfare

Data element concept attributes

Relational attributes

Data Elements
implementing this Data
Element Concept:

[Adult—waist circumference risk indicator, Caucasian adult code N
Health!, Standard 01/03/2005](#)