

Person—lower limb amputation due to vascular disease, code N

Exported from METEOR (AIHW's Metadata Online Registry)

© Australian Institute of Health and Welfare 2024

This product, excluding the AIHW logo, Commonwealth Coat of Arms and any material owned by a third party or protected by a trademark, has been released under a Creative Commons BY 4.0 (CC BY 4.0) licence. Excluded material owned by third parties may include, for example, design and layout, images obtained under licence from third parties and signatures. We have made all reasonable efforts to identify and label material owned by third parties.

You may distribute, remix and build on this website's material but must attribute the AIHW as the copyright holder, in line with our attribution policy. The full terms and conditions of this licence are available at <https://creativecommons.org/licenses/by/4.0/>.

Enquiries relating to copyright should be addressed to info@aihw.gov.au.

Enquiries or comments on the METEOR metadata or download should be directed to the METEOR team at meteor@aihw.gov.au.

Person—lower limb amputation due to vascular disease, code N

Identifying and definitional attributes

Metadata item type:	Data Element
Short name:	Lower limb amputation due to vascular disease
METEOR identifier:	270162
Registration status:	HealthI , Standard 01/03/2005
Definition:	Whether a person has undergone an amputation of toe, forefoot or leg (above or below knee), due to vascular disease, as represented by a code.
Data Element Concept:	Person—lower limb amputation due to vascular disease
Value Domain:	Lower limb amputation due to vascular disease code N

Value domain attributes

Representational attributes

Representation class:	Code
Data type:	Number
Format:	N
Maximum character length:	1

	Value	Meaning
Permissible values:	1	Lower limb amputation - occurred in the last 12 months
	2	Lower limb amputation - occurred prior to the last 12 months
	3	Lower limb amputation - occurred both in and prior to the last 12 months
	4	No history of lower limb amputation due to vascular disease
Supplementary values:	9	Not stated/inadequately described

Collection and usage attributes

Collection methods:	Ask the individual if he/she has had an amputated toe or forefoot or leg (above or below knee), not due to trauma or causes other than vascular disease. If so determine when it was undertaken; within or prior to the last 12 months (or both). Alternatively obtain this information from appropriate documentation.
----------------------------	---

Data element attributes

Source and reference attributes

Submitting organisation:	National Diabetes Data Working Group
Origin:	National Diabetes Outcomes Quality Review Initiative (NDOQRIN) data dictionary.

Reference documents: Duffy MD, John C and Patout MD, Charles A. 1990. Management of the Insensitive Foot in Diabetes: Lessons from Hansen's Disease. *Military Medicine*, 155:575-579. Edmonds M, Boulton A, Buckenham T et al. Report of the Diabetic Foot and Amputation Group. *Diabet Med* 1996; 13: S27-42. Sharon R O'Rourke and Stephen Colagiuri: *The Lower Limb in People With Diabetes*; Content 1997/98 Australian Diabetes Society. Colagiuri S, Colagiuri R, Ward J. *National Diabetes Strategy and Implementation Plan*. Canberra: Diabetes Australia, 1998.

Relational attributes

Related metadata references: Is re-engineered from  [Lower limb amputation due to vascular disease, version 1, DE, NHDD, NHIMG, Superseded 01/03/2005.pdf](#) (17.6 KB)
No registration status

Implementation in Data Set Specifications: [Diabetes \(clinical\) DSS](#)
[Health!](#), Superseded 21/09/2005

DSS specific information:

In people with diabetes, amputations are 15 times more common than in people without diabetes, and 50% of all amputations occur in people with diabetes (*The Lower Limb in People With Diabetes*; 1997/98 Australian Diabetes Society).

Diabetic foot disease is the most common cause of hospitalisation in people with diabetes. Diabetic foot complications are common in the elderly, and amputation rates increase with age: by threefold in those aged 45 - 74 years and sevenfold in population aged over 75 years. As stated by Duffy and authors the rate of lower extremity amputations can be reduced by 50% by the institution of monofilament testing in a preventive care program.

[Diabetes \(clinical\) NBPDS](#)
[Health!](#), Standard 21/09/2005

DSS specific information:

In people with diabetes, amputations are 15 times more common than in people without diabetes, and 50% of all amputations occur in people with diabetes (*The Lower Limb in People With Diabetes*; 1997/98 Australian Diabetes Society).

Diabetic foot disease is the most common cause of hospitalisation in people with diabetes. Diabetic foot complications are common in the elderly, and amputation rates increase with age: by threefold in those aged 45 - 74 years and sevenfold in population aged over 75 years. As stated by Duffy and authors the rate of lower extremity amputations can be reduced by 50% by the institution of monofilament testing in a preventive care program.