Tobacco smoking - consumption/quantity (cigarettes)

Important note: This is an archived metadata standard from the AIHW Knowledgebase. For current metadata standards and related information please access METeOR, the AIHW's Metadata Online Registry at http://meteor.aihw.gov.au

Identifying and Definitional Attributes

Data Dictionary:	NHDD			
Knowledgebase ID:	000403	Version number: 1		
Metadata type:	DATA ELEMENT			
Registration Authority:	NHIMG	Admin status: SUPERSEDED		
		Effective date: 01-MAR-05		
Definition:	The number of cigarettes (manufactured or roll-your-own) smoked per day by a person.			
Context:	 Public health and health care: The number of cigarettes smoked is an important measure of the magnitude of the tobacco problem for an individual. Research shows that of Australians who smoke, the overwhelming majority smoke cigarettes (manufactured or roll-your-own) rather than other tobacco products. From a public health point of view, consumption level is relevant only for regular smokers (those who smoke daily or at least weekly). Data on quantity smoked can be used to: evaluate health promotion and disease prevention programs (assessment of interventions) monitor health risk factors and progress towards National Health Goals and Targets ascertain determinants and consequences of smoking assess a person's exposure to tobacco smoke. 			
Deletional and Depresentational Attributes				

Relational and Representational Attributes

Datatype:	Numeric	
Representational form:	QUANTITAT	TVE VALUE
Representation layout:	NN	
Minimum Size:	1	
Maximum Size:	2	
Data Domain:	99	Not stated/inadequately described

	NOVAL	Count of the number of cigarettes smoked daily.
Guide For Use:	smoke cigare Daily consum consumption consumption number.	nent is relevant only for persons who currently ttes daily or at least weekly. Inption should be reported, rather than weekly Weekly consumption is converted to daily by dividing by 7 and rounding to the nearest whole eater than 98 (extremely rare) should be coded 98.
Collection Methods:	Standard Que interviewer a administered persons aged Available from	nded standard for collecting this information is the estions on the Use of Tobacco Among Adults - dministered (Questions 3a and 3b) and self- (Questions 2a and 2b) versions. The questions cover 18 and over. m the National Centre for Monitoring Cardiovascular e AIHW, telephone (02) 6244 1000.
Related metadata:	is qualified by	y Tobacco smoking - frequency version 1 y Tobacco smoking - product version 1 y Date of birth version 4

Administrative Attributes

Source Document: Standard Questions on the Use of Tobacco Among Adults (1998) Source Organisation: Australian Institute of Health and Welfare

Comments: Where this information is collected by survey and the sample permits, population estimates should be presented by sex and 5-year age groups. Summary statistics may need to be adjusted for age and other relevant variables.

It is recommended that in surveys of smoking, data on age, sex and other socio-demographic variables should be collected.

It is also recommended that when smoking is investigated in relation to health, data on other risk factors including pregnancy status, physical activity, overweight and obesity, and alcohol consumption should be collected.

Data Element Links

Information Model Entities linked to this Data Element

NHIM Lifestyle characteristic

Data Agreements which include this Data Element

DSS - Cardiovascular disease (clinical) From 01-Jan-03 to