# Resuscitation of baby

Important note: This is an archived metadata standard from the AIHW Knowledgebase. For current metadata standards and related information please access METeOR, the AIHW's Metadata Online Registry at <a href="http://meteor.aihw.gov.au">http://meteor.aihw.gov.au</a>

### Identifying and Definitional Attributes

Data Dictionary: NHDD

Knowledgebase ID: 000145 Version number: 2

Metadata type: DATA ELEMENT

Registration NHIMG Admin status: SUPERSEDED

Authority: Effective date: 01-MAR-05

Definition: Active measures taken immediately after birth to establish

independent respiration and heart beat, or to treat depressed

respiratory effort and to correct metabolic disturbances

Context: Perinatal:

Required to analyse need for resuscitation after complications of labour and delivery and to evaluate level of services needed for

different birth settings.

### Relational and Representational Attributes

Datatype: Numeric

Representational

form:

Representation N

layout:

Minimum Size: 1
Maximum Size: 1

Data Domain: 1 None

2 Suction only

3 Oxygen therapy only

4 Intermittent positive pressure respiration (IPPR)

through bag and mask

5 Endotracheal intubation and IPPR

6 External cardiac massage and ventilation

9 Not stated

Guide For Use: This item does not include drug therapy. Code the most severe

measure used. If oxygen is given by bag and mask without IPPR,

code as 'oxygen therapy'.

Related metadata: is used in conjunction with Status of the baby version 1

is used in conjunction with Apgar score version 1

supersedes previous data element Resuscitation of baby version 1 is used in conjunction with Apgar score at 5 minutes version 1

## Administrative Attributes

Source Document:

Source Organisation: National Perinatal Data Development Committee

#### Data Element Links

Information Model Entities linked to this Data Element

NHIM Birth event

Data Agreements which include this Data Element