Person—hypertension treatment with antihypertensive medication indicator (current), code



Exported from METEOR (AIHW's Metadata Online Registry)

© Australian Institute of Health and Welfare 2024

This product, excluding the AIHW logo, Commonwealth Coat of Arms and any material owned by a third party or protected by a trademark, has been released under a Creative Commons BY 4.0 (CC BY 4.0) licence. Excluded material owned by third parties may include, for example, design and layout, images obtained under licence from third parties and signatures. We have made all reasonable efforts to identify and label material owned by third parties.

You may distribute, remix and build on this website's material but must attribute the AIHW as the copyright holder, in line with our attribution policy. The full terms and conditions of this licence are available at https://creativecommons.org/licenses/by/4.0/.

Enquiries relating to copyright should be addressed to info@aihw.gov.au.

Enquiries or comments on the METEOR metadata or download should be directed to the METEOR team at meteor@aihw.gov.au.

Person—hypertension treatment with antihypertensive medication indicator (current), code N

Identifying and definitional attributes

Metadata item type: Data Element

Short name: Hypertension - treatment

METEOR identifier: 302442

Registration status: Health!, Standard 21/09/2005

Definition: Whether a person is currently being treated for hypertension (high blood pressure)

using antihypertensive medication, as represented by a code.

Data element concept attributes

Identifying and definitional attributes

Data element concept: Person—hypertension treatment with antihypertensive medication indicator

METEOR identifier: 304497

Registration status: Health!, Standard 21/09/2005

Definition: Whether an individual is being treated for hypertension (high <u>blood pressure</u>)

using antihypertensive medication.

Object class: Person

Property: Hypertension treatment with antihypertensive medication indicator

Source and reference attributes

Submitting organisation: National diabetes data working group

Value domain attributes

Identifying and definitional attributes

Value domain: Yes/no/not stated/inadequately described code N

METEOR identifier: 301747

Registration status: Health!, Standard 21/09/2005

Housing assistance, Standard 10/02/2006

Community Services (retired), Standard 14/02/2006

Early Childhood, Standard 21/05/2010 Homelessness, Standard 23/08/2010

Independent Hospital Pricing Authority, Standard 01/11/2012

<u>Disability</u>, Standard 07/10/2014 <u>Indigenous</u>, Standard 13/03/2015

Children and Families, Standard 22/11/2016

Definition: A code set representing 'yes', 'no' and 'not stated/inadequately described'.

Representational attributes

Representation class: Code

Data type: Number

Format: N

Maximum character length: 1

Value Meaning

Permissible values: 1 Yes

2 No

Supplementary values: 9 Not stated/inadequately described

Collection and usage attributes

Guide for use: CODE 9 Not stated/inadequately described

This code is not for use in primary data collections.

Data element attributes

Collection and usage attributes

Guide for use: CODE 1 Yes

Record if a person is currently being treated for hypertension using antihypertensive

medication.

CODE 2 No

Record if a person is not currently being treated for hypertension using

antihypertensive medication.

Collection methods: Ask the individual if he/she is currently treated with anti-hypertensive medications.

Alternatively obtain the relevant information from appropriate documentation.

Source and reference attributes

Submitting organisation: National diabetes data working group

Origin: National Diabetes Outcomes Quality Review Initiative (NDOQRIN) data dictionary.

Reference documents: Pahor M, Psaty BM, Furberg CD. Treatment of hypertensive patients with diabetes.

Lancet 1998; 351:689-90. Tight blood pressure control and risk of macrovascular and microvascular complications in type 2 diabetes: UKPDS 38. UK Prospective Diabetes Study Group [erratum appears in Br Med J 1999; 318:29]. Br Med J

1998; 317:703-13.

Grossman E, Messerli FH, Goldbourt U, Curb JD, Pressel SL, Cutler JA, Savage PJ, Applegate WB, Black H, et al. Effect of diuretic-based antihypertensive treatment on cardiovascular disease risk in older diabetic patients with isolated systolic hypertension. Systolic Hypertension in the Elderly Program Cooperative

Research Group. JAMA 1996; 276:1886-92.

Hypertension in diabetes [Australian Prescriber Feb 2002]. American Journal of

Preventive Medicine 2002;21.

Relational attributes

Related metadata references:

Supersedes Person—hypertension treatment status (antihypertensive medication),

code N

Health!, Superseded 21/09/2005

Specifications:

Health!, Superseded 01/09/2012

> Acute coronary syndrome (clinical) DSS Health!, Superseded 02/05/2013

Acute coronary syndrome (clinical) NBPDS 2013-

Health!, Standard 02/05/2013 Implementation start date: 01/07/2013

Diabetes (clinical) NBPDS Health!, Standard 21/09/2005

DSS specific information:

Hypertension is probably the most important public health problem in developed countries. It is common, asymptomatic, readily detectable, usually easily treatable, and often leads to lethal complications if left untreated.

Elevated blood pressure (Hypertension) is a recognised risk for microvascular and macro vascular complications of diabetes (coronary, cerebral and peripheral).

Hypertension is elevated arterial blood pressure above the normal range (130 to 139/85 to 89 mm Hg) and values above these are defined as hypertension. Lower levels of target blood pressure should be aimed for in specific groups, e.g. in diabetics aim for blood pressure less than 135/80 mm Hg.

Many diabetics fail to control high blood pressure. Among all the diabetics with high blood pressure, 29% were unaware that they had high blood pressure and only slightly more than half were receiving hypertensive medications as treatment. Numbers of studies have shown that good management of blood pressure is at least as important as good control of blood glucose and the reduction of cholesterol in preventing the complications of diabetes.

Antihypertensives - Australian Medicines Handbook: February, 2001. Tight blood control in diabetes usually requires combination therapy as stated by (Australian Diabetes society) Therapeutic Guidelines Limited (05.04.2002).

People taking antihypertensives are also encouraged to make healthy lifestyle changes, such as guit smoking, lose weight and have regular physical activity. The level of blood pressure should generally be established on at least two to four occasions prior to initiating antihypertensive medication.

Systematic reviews of studies that have reported outcomes in patients with diabetes and hypertension indicate that combination therapy is frequently required and may be more beneficial than monotherapy. In the past multi-drug therapy to control hypertension has not been advocated much, but according to the special report published in the American Journal of Kidney Diseases, if ACE inhibitor therapy alone doesn't achieve good blood pressure control, multi-drug therapy should be implemented. (Heart Center Online)