© Australian Institute of Health and Welfare 2024

This product, excluding the AIHW logo, Commonwealth Coat of Arms and any material owned by a third party or protected by a trademark, has been released under a Creative Commons BY 4.0 (CC BY 4.0) licence. Excluded material owned by third parties may include, for example, design and layout, images obtained under licence from third parties and signatures. We have made all reasonable efforts to identify and label material owned by third parties.

You may distribute, remix and build on this website's material but must attribute the AlHW as the copyright holder, in line with our attribution policy. The full terms and conditions of this licence are available at https://creativecommons.org/licenses/by/4.0/.

Enquiries relating to copyright should be addressed to info@aihw.gov.au.

Enquiries or comments on the METEOR metadata or download should be directed to the METEOR team at meteor@aihw.gov.au.

National Healthcare Agreement: Pl 16-Potentially avoidable deaths, 2013

Identifying and definitional attributes

Metadata item type: Indicator

Indicator type: Progress measure

Short name: PI 16-Potentially avoidable deaths, 2013

METEOR identifier: 497242

Registration status: Health!, Superseded 30/04/2014

Description: Deaths that are potentially avoidable within the present health system:

· potentially preventable deaths are those amenable to screening and primary

prevention, such as immunisation

· deaths from potentially treatable conditions are those amenable to therapeutic

interventions

Indicator set: National Healthcare Agreement (2013)

Health!, Superseded 30/04/2014

Outcome area: Primary and Community Health

Health!, Standard 07/07/2010

Data quality statement: National Healthcare Agreement: PI 16-Potentially avoidable deaths, 2013 QS

Health!, Superseded 14/01/2015

Collection and usage attributes

Population group age to: 75 years

Computation description: Deaths are defined as avoidable in the context of the present health system.

International Classification of Diseases (ICD-10) codes in scope are as specified

below:

Cause of death group	ICD-10 Codes	Limits (age, sex)	Treatable	Preventable
Infections				
Tuberculosis	A15-A19, B90		X	
Selected invasive bacterial and protozoal infections	A38-A41, A46, A48.1 B50-B54, G00,G03, J02.0, J13-J15, J18, L03		X	
Hepatitis	B15–B19			Χ
HIV/AIDS	B20-B24			Χ
Viral pneumonia and influenza	J10, J12, J21			X
Enteritis and other diarrhoeal disease	A00-A09			X
Childhood vaccine- preventable diseases	A35–A37, A49.1, A49.2, A80, B01, B05–B06, J11	0–14		X

Cause of death group	ICD-10 Codes	Limits (age, sex)	Treatable	Preventable
Sexually transmitted diseases, except HIV/AIDS	A50-A64, M02.3, N34.1, N70-N73, N75.0, N75.1, N76.4, N76.6, O00			X
Ear infections—otitis media and mastoiditis	H65–H70		X	
Upper respiratory tract infection	J00–J06, except for J02.0		X	
Cancer				
Lip, oral cavity and pharynx	C00-C14			X
Oesophagus	C15			X
Stomach	C16			Χ
Colorectal	C18-C21		X	
Liver	C22			Χ
Lung	C33, C34			Χ
Melanoma of skin	C43		Χ	
Non-melanocytic skin	C44		Χ	
Breast	C50	Female	Χ	
Cervix	C53		Χ	
Uterus	C54, C55		Χ	
Bladder	C67		X	
Thyroid	C73		X	
Hodgkin's disease	C81		X	
Lymphoid leukaemia— acute/chronic	C91.0, C91.1		X	
Benign cancers	D10-D36		Χ	
Malignant neoplasm of testis	C62			X
Malignant neoplasm of eye and adnexa	C69			X
Diabetes	E10-E14		X (0.5)	X (0.5)
Drug use disorders				
Alcohol related disease	F10, I42.6, K29.2, K70			X
Illicit drug use disorders	F11–F16, F18, F19			X
Epilepsy	G40, G41		X	
Diseases of the circulatory system				
Rheumatic and other valvular heart disease	101–109		X	
Hypertensive heart disease	l11		X	
Ischaemic heart disease	120–125		X (0.5)	X (0.5)

Cause of death group	ICD-10 Codes	Limits (age, sex)	Treatable	Preventable
Cerebrovascular diseases	160–169		X (0.5)	X (0.5)
Aortic aneurysm	l71			X
Phlebitis and thrombophlebitis of other deep vessels of lower extremities	180.2			X
Pulmonary embolism	126			X
Diseases of the genitourinary system				
Nephritis and nephrosis	I12, I13, N00-N07, N17-N19		X	
Obstructive uropathy & prostatic hyperplasia	N13, N20, N21, N35, N40		X	
Diseases of the respiratory system				
COPD	J40-J44	45–74 years		X
Asthma	J45, J46	0–44 years	Χ	
Diseases of the digestive system				
Peptic ulcer disease	K25-K28		Χ	
Diseases of appendix; hernia; disorders of gallbladder, biliary tract, and pancreas	K35–K38, K40– K46, K80–K83, K85, K86		X	
Chronic liver disease (excluding alcohol- related disease)	K73, K74			X
Maternal & infant causes				
Birth defects	H31.1, P00, P04, Q00–Q99		X	
Complications of perinatal period	P03, P05–P95		Χ	
Unintentional injuries				
Road traffic injuries	V01–V04, V06, V09–V80, V87, V89, V99			Х
Falls	W00-W19			Χ
Fires, burns	X00-X09			Х
Accidental poisonings	X40-X49			X
Drownings	W65-W74			Х
War	Y36			X
Intentional injuries				
Suicide and self- inflicted injuries	X60-X84, Y87.0, Y10-Y34			X

Cause of death group	ICD-10 Codes	Limits (age, sex)	Treatable	Preventable
Violence	X85-Y09, Y87.1			Χ
Other conditions				
Nutritional deficiency anaemia	D50-D53			X
Thyroid disorders	E00-E07		Χ	
Adrenal disorders	E24, E27		Χ	
Congenital metabolic disorders	E25, E70.0, E74.2		X	
Osteomyelitis and other osteopathies of bone	M86, M89			X
Complications of pregnancy, labour or the puerperium	O01–O99			X
Misadventures to patients during surgical and medical care	Y60-Y69			X
Medical devices associated with adverse incidents in diagnostic & therapeutic use	Y70–Y82			X
Surgical and other medical procedures as the cause of abnormal reaction of the patient, or of later complication, without mention of misadventure at the time of the procedure	Y83-Y84			X

Rates are directly age-standardised to the 2001 Australian population.

Variability bands are to be calculated for single-year rates using the method below.

Presented per 100,000 population.

Computation: 100,000 x (Numerator ÷ Denominator)

Calculated separately for 'preventable' and 'treatable' categories.

Variability bands are to be calculated for single-year rates using the following method for estimating 95% confidence intervals:

Age-standardised rate

$$CI(ASR)_{SSN_t} = ASR \pm 1.96 \times \sqrt{\sum_{t=1}^{I} \frac{w_t^2 d_t}{n_t^2}}$$

Where w = the proportion of the standard population in age group i

d=the number of deaths in age group i

n=the number of people in the population in age group i

Numerator: Number of deaths of persons aged less than 75 categorised as potentially

avoidable

Numerator data elements:

Data Element / Data Set

Person—underlying cause of death, code (ICD-10 2nd edn) ANN-ANN

Data Source

ABS Causes of Death Collection

Guide for use

Data source type: Registry

Denominator:

Population aged less than 75 years

Denominator data elements:

-Data Element / Data Set-

Person—estimated resident population of Australia

Data Source

ABS Estimated resident population (total population)

Guide for use

Data source type: Census-based plus administrative by-product data

Data Element / Data Set-

Person—estimated resident population of Australia

Data Source

ABS Indigenous experimental estimates and projections (2001 Censusbased)

Guide for use

Data source type: Census-based plus administrative by-product data

Data Element / Data Set-

Person—age, total years N[NN]

Data Source

ABS Estimated resident population (total population)

Guide for use

Data source type: Census-based plus administrative by-product data

Data Element / Data Set-

Person—age, total years N[NN]

Data Source

ABS Indigenous experimental estimates and projections (2001 Censusbased)

Guide for use

Data source type: Census-based plus administrative by-product data

Disaggregation:

2007, 2008, 2009 (resupplied for revision to ABS WA and national Indigenous deaths data), 2010—State and territory, by preventable and treatable categories.

2007, 2008, 2009 (resupplied for revision to ABS WA and national Indigenous death data), 2010—Nationally, by preventable and treatable categories:

Indigenous status

2006–2010—State and territory, by preventable and treatable categories, by:

Indigenous status

Some disaggregations may result in numbers too small for publication. Disaggregation by Indigenous status will be based on data only from jurisdictions for which the quality of Indigenous identification is considered acceptable—NSW, Qld, SA, WA, NT.

Disaggregation data elements:

Data Element / Data Set

Person—Indigenous status, code N

Data Source

ABS Causes of Death Collection

Guide for use

Data source type: Registry

-Data Element / Data Set-

Person—area of usual residence, geographical location code (ASGC 2007) NNNNN

Data Source

ABS Causes of Death Collection

Guide for use

Data source type: Registry

Used for disaggregation by state/territory

Comments:

Most recent data available for 2013 Council of Australian Governments (COAG) Reform Council (CRC) report:

- 2010 (Total population)
- Aggregated data 2006–2010 (Indigenous status)

Due to small number of Indigenous deaths reported each year, 5 year combined data will be reported for the current reporting period.

A system error affected the reporting of WA Aboriginal and Torres Strait Islander death registrations for 2007, 2008 and 2009 which affected the WA and national death rates reported for the 2011 and 2012 CRC reports. These data have been revised for 2013 reporting.

Data by remoteness may be available, pending assessment of data quality.

Representational attributes

Representation class: Rate

Data type: Real

Unit of measure: Person

Format: NN[N].N

Indicator conceptual framework

Framework and dimensions:

Deaths

Data source attributes

Data sources:

Data Source

ABS Estimated resident population (total population)

Frequency

Quarterly

Data quality statement

ABS Estimated resident population (total population), QS

Data custodian

Australian Bureau of Statistics

Data Source

ABS Causes of Death Collection

Frequency

Annual

Data quality statement

ABS causes of death collection, QS

Data custodian

Australian Bureau of Statistics

Data Source

ABS Indigenous experimental estimates and projections (2001 Censusbased)

Frequency

Periodic

Data quality statement

ABS Indigenous experimental estimates and projections, QS

Data custodian

Australian Bureau of Statistics

Accountability attributes

Reporting requirements: National Healthcare Agreement

Organisation responsible

for providing data:

Australian Bureau of Statistics (ABS).

collection required:

Further data development / Specification: Minor work required, the measure needs minor work to meet the

intention of the indicator.

Relational attributes

Related metadata references:

Supersedes National Healthcare Agreement: PI 20-Potentially avoidable deaths, 2012

Health!, Superseded 25/06/2013

Has been superseded by <u>National Healthcare Agreement</u>: <u>PI 16-Potentially avoidable deaths</u>, 2014

Health!, Superseded 14/01/2015

See also National Healthcare Agreement: PI 03-Prevalence of overweight and obesity, 2013

Health!, Superseded 30/04/2014

See also National Healthcare Agreement: PI 04-Rates of current daily smokers, 2013

Health!, Superseded 30/04/2014

See also National Healthcare Agreement: PI 05-Levels of risky alcohol consumption, 2013

Health!, Superseded 30/04/2014

See also National Healthcare Agreement: PI 06-Life expectancy, 2013 Health!, Superseded 30/04/2014

See also National Healthcare Agreement: PI 07-Infant and young child mortality rate, 2013

Health!, Superseded 30/04/2014

See also National Healthcare Agreement: PI 08-Major causes of death, 2013 Health!, Superseded 30/04/2014

See also National Healthcare Agreement: PI 18-Selected potentially preventable hospitalisations, 2013

Health!, Superseded 30/04/2014

See also National Healthcare Agreement: PI 23-Unplanned hospital readmission rates, 2013

Health!, Superseded 30/04/2014