

# National Healthcare Agreement: PI 16-Potentially avoidable deaths, 2014 QS

Exported from METEOR (AIHW's Metadata Online Registry)

© Australian Institute of Health and Welfare 2024

This product, excluding the AIHW logo, Commonwealth Coat of Arms and any material owned by a third party or protected by a trademark, has been released under a Creative Commons BY 4.0 (CC BY 4.0) licence. Excluded material owned by third parties may include, for example, design and layout, images obtained under licence from third parties and signatures. We have made all reasonable efforts to identify and label material owned by third parties.

You may distribute, remix and build on this website's material but must attribute the AIHW as the copyright holder, in line with our attribution policy. The full terms and conditions of this licence are available at <https://creativecommons.org/licenses/by/4.0/>.

Enquiries relating to copyright should be addressed to [info@aihw.gov.au](mailto:info@aihw.gov.au).

Enquiries or comments on the METEOR metadata or download should be directed to the METEOR team at [meteor@aihw.gov.au](mailto:meteor@aihw.gov.au).

# National Healthcare Agreement: PI 16-Potentially avoidable deaths, 2014 QS

## Identifying and definitional attributes

<b>Metadata item type:</b>	Data Quality Statement
<b>METEOR identifier:</b>	517743
<b>Registration status:</b>	<a href="#">Health!</a> , Superseded 14/01/2015

## Data quality

**Institutional environment:** These collections are conducted under the *Census and Statistics Act 1905*. For information on the institutional environment of the ABS, including the legislative obligations of the Australian Bureau of Statistics (ABS), financing and governance arrangements, and mechanisms for scrutiny of ABS operations, see [ABS Institutional Environment](#).

**Timeliness:** Causes of death data is published on an annual basis. Death records are provided electronically to the ABS by individual Registrars on a monthly basis for compilation into aggregate statistics on a quarterly and annual basis. One dimension of timeliness in death registrations data is the interval between the occurrence and registration of a death. As a result, a small number of deaths occurring in one year are not registered until the following year or later.

Preliminary Estimated Resident Population (ERP) data is compiled and published quarterly and is generally made available five to six months after the end of each reference quarter.

Every year, the 30 June ERP is further disaggregated by sex and single year of age, and is made available five to six months after the end of the reference quarter. Commencing with data for September quarter 2006, revised estimates are released annually and made available 21 months after the end of the reference period for the previous financial year, once more accurate births, deaths and net overseas migration data becomes available. In the case of births and deaths, the revised data is compiled on a date of occurrence basis. In the case of net overseas migration, final data is based on actual traveller behaviour. Generally ERP data is not changed once it has been finalised unless there are compelling reasons to do so, as in June 2013 when data from September 1991 to June 2006 was revised (for more information on this recasting process, please see the feature article titled *Recasting 20 years of ERP in the December quarter 2012 issue of Australian Demographic Statistics* (cat. no. 3101.0).

For further information on ABS Estimated Resident Population, see the relevant Data Quality Statement.

**Accessibility:** Data for this indicator have been age-standardised, using the direct method, to 'under 75 years' of age. Direct age-standardisation to the 2001 total Australian population was used (see *Data Cube: Standard Population for Use in Age-Standardisation Table in Australian Demographic Statistics*, Dec 2012 (cat. no. 3101.0)). Age-standardised results provide a measure of relative difference only between populations.

**Interpretability:** Causes of death data are available in a variety of formats on the ABS website under the 3303.0 product family. ERP data is available in a variety of formats on the ABS website under the 3101.0 and 3201.0 product families. Further information on deaths and mortality may be available on request. The ABS observes strict confidentiality protocols as required by the *Census and Statistics Act (1905)*. This may restrict access to data at a very detailed level.

**Relevance:**

The ABS Causes of Death collection includes all deaths that occurred and were registered in Australia, including deaths of persons whose usual residence is overseas. Deaths of Australian residents that occurred outside Australia may be registered by individual Registrars, but are not included in ABS deaths or causes of death statistics.

Data in the Causes of Death collection include demographic items, as well as causes of death information, which is coded according to the International Statistical Classification of Diseases and Related health Problems (ICD). ICD is the international standard classification for epidemiological purposes and is designed to promote international comparability in the collection, processing, classification, and presentation of causes of death statistics. The classification is used to classify diseases and causes of disease or injury as recorded on many types of medical records as well as death records. The ICD has been revised periodically to incorporate changes in the medical field. The 10th revision of ICD (ICD-10) has been used by the ABS to code cause of death since 1997.

For further information on the ABS Causes of Death collection, see the relevant Data Quality Statement.

**Accuracy:**

Information on causes of death is obtained from a complete enumeration of deaths registered during a specified period and is not subject to sampling error. However, deaths data sources are subject to non-sampling error which can arise from inaccuracies in collecting, recording and processing the data.

Although it is considered likely that most deaths of Aboriginal and Torres Strait Islander (Indigenous) Australians are registered, a proportion of these deaths are not registered as Indigenous. Information about the deceased is supplied by a relative or other person acquainted with the deceased, or by an official of the institution where the death occurred and may differ from the self-identified Indigenous origin of the deceased. Forms are often not subject to the same best practice design principles as statistical questionnaires, and respondent and/or interviewer understanding is rarely tested. Over-precise analysis of Indigenous deaths and mortality should be avoided.

All coroner certified deaths registered after 1 January 2006 are now subject to a revisions process. In this round of National Healthcare Agreement (NHA) reporting, 2007, 2008 and 2009 data is final, 2010 data is revised and 2011 data is preliminary. Data for 2010 and 2011 is subject to further revisions. Prior to 2006 all ABS processing of causes of death data for a particular reference period was finalised approximately 13 months after the end of the reference period. Where insufficient information was available to code a cause of death (e.g. a coroner certified death was yet to be finalised by the Coroner), less specific ICD codes were assigned as required by the ICD coding rules. The revision process enables the use of additional information relating to coroner certified deaths, as it becomes available over time. This results in increased specificity of the assigned ICD-10 codes.

Revisions will only impact on coroner certified deaths, as further information becomes available to the ABS about the causes of these deaths. See Technical Note: Causes of Death Revisions 2009 and 2010 in Causes of Death, Australia, 2011 (cat.no. 3303.0).

In November 2010, the Queensland Registrar of Births, Deaths and Marriages advised the ABS of an outstanding deaths registration initiative undertaken by the Registry. This initiative resulted in the November 2010 registration of 374 previously unregistered deaths which occurred between 1992 and 2006 (including a few for which a date of death was unknown). Of these, around three-quarters (284) were deaths of Aboriginal and Torres Strait Islander Australians.

The ABS discussed different methods of adjustment of Queensland death registrations data for 2010 with key stakeholders. Following the discussion, a decision was made by the ABS and key stakeholders to use an adjustment method that added together deaths registered in 2010 for usual residents of Queensland which occurred in 2007, 2008, 2009 and 2010. This method minimises the impact on mortality indicators used in various government reports. However, care should still be taken when interpreting Aboriginal and Torres Strait Islander death data for Queensland for 2010. Please note that there are differences between data output in the Causes of Death, Australia, 2010 publication (cat. no. 3303.0) and 2010 data reported for the NHA, as this adjustment was not applied in the publication. For

further details see Technical Note: Registration of outstanding deaths, Queensland 2010, from the Deaths, Australia, 2010 publication (cat. no. 3302.0) and Explanatory Note 103 in the Causes of Death, Australia, 2010 publication (cat. no. 3303.0).

Investigation conducted by the WA Registrar of Births, Deaths and Marriages indicated that some deaths of non-Indigenous people were wrongly identified as deaths of Indigenous people in WA for 2007, 2008 and 2009. The ABS discussed this issue with a range of key stakeholders and users of Aboriginal and Torres Strait Islander deaths statistics. Following this discussion, the ABS did not release WA Aboriginal and Torres Strait Islander deaths data for the years 2007, 2008 and 2009 in the 2010 issue of Deaths, Australia publication, or in the 2011 NHA data supply. The WA Registry corrected the data and resupplied the corrected data to the ABS. These corrected data were then released by the ABS in spreadsheets attached to Deaths, Australia, 2010 (cat. no. 3302.0) publication on 24 May 2012, and are included in this round of NHA reporting.

All ERP data sources are subject to non-sampling error. Non-sampling error can arise from inaccuracies in collecting, recording and processing the data. In the case of Census and Post Enumeration Survey (PES) data, every effort is made to minimise reporting error by the careful design of questionnaires, intensive training and supervision of interviewers, and efficient data processing procedures. The ABS does not have control over any non-sampling error associated with births, deaths and migration data. For more information see the Demography Working Paper 1998/2 - Quarterly birth and death estimates, 1998 (cat. no. 3114.0). and Australian Demographic Statistics (cat. no. 3101.0).

Non-Indigenous estimates are available for census years only. In the intervening years, Indigenous population projections are based on assumptions about past and future levels of fertility, mortality and migration. In the absence of non-Indigenous population figures for these years, it is possible to derive denominators for calculating non-Indigenous rates by subtracting the projected Indigenous population from the total population. For the current round of NHA reporting, in the absence of 2011 Census-based Indigenous population projections, the non-Indigenous population denominator has been calculated by subtracting the 2006 Census-based Indigenous projections (see Experimental Estimates and Projections, Aboriginal and Torres Strait Islander Australians, August 2009, cat. no. 3238.0) from the 2006 Census-based Estimated Resident Population (3101.0). Such figures have a degree of uncertainty and should be used with caution, particularly as the time from the base year of the projection series increases.

Non-Indigenous data from the Causes of Death collection do not include death registrations with a 'not stated' Indigenous status.

Some rates are unreliable due to small numbers of deaths over the reference period. Resultant rates could be misleading, for example, where the non-Indigenous mortality rate is higher than the indigenous mortality rate. Age-standardised death rates based on a very low death count have been deemed unpublishable. Some cells have also not been published to prevent back-calculation of these suppressed cells. Caution should be used when interpreting rates for this indicator.

**Coherence:**

The methods used to construct the indicator are consistent and comparable with other collections and with international practice.

**Relational attributes**

**Related metadata references:**

Supersedes [National Healthcare Agreement: PI 16-Potentially avoidable deaths, 2013 QS](#)

[Health!](#), Superseded 14/01/2015

Has been superseded by [National Healthcare Agreement: PI 16-Potentially avoidable deaths, 2015 QS](#)

[Health!](#), Superseded 08/07/2016

**Indicators linked to this Data Quality statement:**

[National Healthcare Agreement: PI 16-Potentially avoidable deaths, 2014](#)

[Health!](#), Superseded 14/01/2015