

Acute coronary syndromes: 6a-ACS patients with an individualised care plan

Exported from METEOR (AIHW's Metadata Online Registry)

© Australian Institute of Health and Welfare 2024

This product, excluding the AIHW logo, Commonwealth Coat of Arms and any material owned by a third party or protected by a trademark, has been released under a Creative Commons BY 4.0 (CC BY 4.0) licence. Excluded material owned by third parties may include, for example, design and layout, images obtained under licence from third parties and signatures. We have made all reasonable efforts to identify and label material owned by third parties.

You may distribute, remix and build on this website's material but must attribute the AIHW as the copyright holder, in line with our attribution policy. The full terms and conditions of this licence are available at <https://creativecommons.org/licenses/by/4.0/>.

Enquiries relating to copyright should be addressed to info@aihw.gov.au.

Enquiries or comments on the METEOR metadata or download should be directed to the METEOR team at meteor@aihw.gov.au.

Acute coronary syndromes: 6a-ACS patients with an individualised care plan

Identifying and definitional attributes

Metadata item type:	Indicator
Indicator type:	Indicator
Short name:	Indicator 6a-ACS patients with an individualised care plan
METEOR identifier:	612083
Registration status:	HealthI , Standard 12/09/2016
Description:	Proportion of acute coronary syndrome (ACS) patients provided with a written, individualised care plan (addressing factors such as gradual physical activity, smoking cessation and therapies addressing psychosocial needs).
Rationale:	ACS patients can successfully contribute to managing their own condition.
Indicator set:	Clinical care standard indicators: acute coronary syndromes HealthI , Standard 12/09/2016
Outcome area:	Individualised care plan HealthI , Standard 12/09/2016

Collection and usage attributes

Computation description:	<p>Both the numerator and the denominator include patients with acute chest pain, or other symptoms suggestive of ACS.</p> <p>Both the numerator and the denominator exclude patients separated on a palliative care pathway, or where adherence to a secondary prevention plan is not indicated.</p> <p>For an individualised care plan to be included for a patient in the numerator, it needs to be a written plan, addressing factors specific to the patient, such as:</p> <ul style="list-style-type: none">• gradual physical activity (except for patients who are very debilitated)• attention to lifestyle factors such as smoking cessation, nutrition, alcohol moderation, and weight management, as appropriate• rest• therapies addressing psychosocial needs• self-administration of short-acting nitrates• self-administration of aspirin; unless contraindicated (most patients should already be taking aspirin)• calling an ambulance (e.g. dialling 000) if chest pain or discomfort is not completely relieved within 10 minutes• individualised clinician notification and care plan for those living in areas where an ambulance is not readily available (Acute Coronary Syndrome Guidelines Working Group 2006). <p>Presented as a percentage.</p>
Computation:	$(\text{Numerator} \div \text{denominator}) \times 100$
Numerator:	Number of ACS patients provided with an individualised care plan.
Denominator:	Number of ACS patients separated from hospital.
Comments:	Resources provided to patients who do not speak English may require translation into other languages.

Representational attributes

Representation class:	Percentage
Data type:	Real

Unit of measure: Episode

Format: N[NN]

Source and reference attributes

Submitting organisation: Australian Commission on Safety and Quality in Health Care

Reference documents: Acute Coronary Syndrome Guidelines Working Group 2006. Guidelines for the management of acute coronary syndromes 2006. Medical Journal of Australia 184(8):S1–S30.