

Pain management

Exported from METEOR (AIHW's Metadata Online Registry)

© Australian Institute of Health and Welfare 2024

This product, excluding the AIHW logo, Commonwealth Coat of Arms and any material owned by a third party or protected by a trademark, has been released under a Creative Commons BY 4.0 (CC BY 4.0) licence. Excluded material owned by third parties may include, for example, design and layout, images obtained under licence from third parties and signatures. We have made all reasonable efforts to identify and label material owned by third parties.

You may distribute, remix and build on this website's material but must attribute the AIHW as the copyright holder, in line with our attribution policy. The full terms and conditions of this licence are available at <https://creativecommons.org/licenses/by/4.0/>.

Enquiries relating to copyright should be addressed to info@aihw.gov.au.

Enquiries or comments on the METEOR metadata or download should be directed to the METEOR team at meteor@aihw.gov.au.

Pain management

Identifying and definitional attributes

Metadata item type:	Outcome Area
METEOR identifier:	628050
Registration status:	Health! , Standard 12/09/2016
Description:	A patient with a hip fracture is assessed for pain at the time of presentation and regularly throughout their hospital stay, and receives pain management including the use of multimodal analgesia, if clinically appropriate.

Relational attributes

Indicator sets linked to this outcome area:	Clinical care standard indicators: hip fracture Australian Commission on Safety and Quality in Health Care , Superseded 18/06/2018 Health! , Standard 12/09/2016 Clinical care standard indicators: hip fracture 2018 Australian Commission on Safety and Quality in Health Care , Standard 15/05/2018
--	--

Indicators linked to this outcome area:	Hip fracture care clinical care standard indicators: 2a-Evidence of local arrangements for timely and effective pain management for hip fracture Health! , Standard 12/09/2016 Hip fracture care clinical care standard indicators: 2a-Evidence of local arrangements for timely and effective pain management for hip fracture Health! , Standard 12/09/2016 Hip fracture care clinical care standard indicators: 2b-Proportion of hip fracture patients who have documented pain assessment within 30 minutes of emergency department presentation and either receive analgesia within this time or do not require it Health! , Standard 12/09/2016 Hip fracture care clinical care standard indicators: 2b-Proportion of hip fracture patients who have documented pain assessment within 30 minutes of emergency department presentation and either receive analgesia within this time or do not require it Health! , Standard 12/09/2016
--	--

Source and reference attributes

Submitting organisation:	Australian Commission on Safety and Quality in Health Care
---------------------------------	--