National Healthcare Agreement: PI 05-Levels of risky alcohol consumption. 2017

Exported from METEOR (AIHW's Metadata Online Registry)
© Australian Institute of Health and Welfare 2024
This product, excluding the AlHW logo, Commonwealth Coat of Arms and any material owned by a third party or protected by a trademark, has been released under a Creative Commons BY 4.0 (CC BY 4.0) licence. Excluded material owned by third parties may include, for example, design and layout, images obtained under licence from third parties and signatures. We have

made all reasonable efforts to identify and label material owned by third parties.

You may distribute, remix and build on this website's material but must attribute the AlHW as the copyright holder, in line with our attribution policy. The full terms and conditions of this licence are available at https://creativecommons.org/licenses/by/4.0/.

Enquiries relating to copyright should be addressed to info@aihw.gov.au.

Enquiries or comments on the METEOR metadata or download should be directed to the METEOR team at meteor@aihw.gov.au.

National Healthcare Agreement: Pl 05-Levels of risky alcohol consumption, 2017

Identifying and definitional attributes

Metadata item type: Indicator

Indicator type: Progress measure

Short name: PI 05–Levels of risky alcohol consumption, 2017

METEOR identifier: 629999

Registration status: Health!, Superseded 30/01/2018

Description: Proportion of adults at risk of long-term harm from alcohol.

Indicator set: National Healthcare Agreement (2017)

Health!, Superseded 30/01/2018

Outcome area: Prevention

Health!, Standard 07/07/2010

Data quality statement: National Healthcare Agreement: PI 05-Levels of risky alcohol consumption, 2017

QS

Health!, Standard 31/01/2017

Collection and usage attributes

Population group age

from:

18 years

Computation description: Data on persons 'at risk of long term alcohol related harm' is based on the 2009

National Health and Medical Research Council (NHMRC) guideline 'for healthy men and women, drinking no more than two standard drinks on any day reduces the lifetime risk of harm from alcohol-related disease or injury' and has been operationalised as: for both males and females, an average of more than 2

standard drinks per day in the last week.

Rates are directly age-standardised to the 2001 Australian population.

Analysis by remoteness and Socio-Economic Indexes for Areas (SEIFA) Index of Relative Socio-economic Disadvantage (IRSD) is based on usual residence of

person.

Presented as a percentage.

95% confidence intervals and relative standard errors are calculated for rates.

Computation: 100 × (Numerator ÷ Denominator)

Numerator: Number of persons aged 18 years or over assessed as having an alcohol

consumption pattern that puts them at risk of long term alcohol-related harm.

Numerator data elements: Data Element / Data Set

Person-age

Data Source

ABS 2014-15 National Health Survey (NHS)

Guide for use

Data source type: Survey

Data Element / Data Set

Person—alcohol consumption amount

Data Source

ABS 2014-15 National Health Survey (NHS)

Guide for use

Data source type: Survey

Data Element / Data Set-

Person—alcohol consumption frequency

Data Source

ABS 2014-15 National Health Survey (NHS)

Guide for use

Data source type: Survey

-Data Element / Data Set-

Person—age

Data Source

ABS 2014-15 National Aboriginal and Torres Strait Islander Social Survey (NATSISS)

Guide for use

Data source type: Survey

-Data Element / Data Set-

Person—alcohol consumption amount

Data Source

ABS 2014-15 National Aboriginal and Torres Strait Islander Social Survey (NATSISS)

Guide for use

Data source type: Survey

Denominator:

Population aged 18 years or over.

Denominator data elements:

Data Element / Data Set-

Person-age

Data Source

ABS 2014-15 National Health Survey (NHS)

Guide for use

Data source type: Survey

-Data Element / Data Set-

Person—age

Data Source

ABS 2014-15 National Aboriginal and Torres Strait Islander Social Survey (NATSISS)

Guide for use

Data source type: Survey

Disaggregation:

2014-15-State and territory, by:

- sex by age (not reported)
- Indigenous status (non-Indigenous not reported)
- remoteness (Australian Statistical Geography Standard (ASGS) Remoteness Structure)
- 2011 SEIFA IRSD quintiles
- · disability status (not reported)

2014–15—Nationally, by (all not reported):

- sex by remoteness (ASGS Remoteness Structure).
- 2011 SEIFA IRSD deciles.
- remoteness (ASGS Remoteness Structure) by 2011 SEIFA IRSD deciles.

Some disaggregation may result in numbers too small for publication.

Disaggregation data elements:

- Data Element / Data Set-

Person-age

Data Source

ABS 2014-15 National Health Survey (NHS)

Guide for use

Data source type: Survey

Data Element / Data Set-

Person—area of usual residence

Data Source

ABS 2014-15 National Health Survey (NHS)

Guide for use

Data source type: Survey

Data Element / Data Set

Person—disability status

Data Source

ABS 2014-15 National Health Survey (NHS)

Guide for use

Data source type: Survey

Data Element / Data Set-

Person—Indigenous status

Data Source

ABS 2014-15 National Health Survey (NHS)

Guide for use

Data source type: Survey

Data Element / Data Set-

Person—area of usual residence

Data Source

ABS 2014-15 National Aboriginal and Torres Strait Islander Social Survey (NATSISS)

Guide for use

Data source type: Survey

Used for disaggregation by state/territory

Comments:

Most recent data available for 2017 National Healthcare Agreement performance reporting: 2014–15 (total population, non-Indigenous: NHS); 2014–15 (Indigenous only: NATSISS).

National Aboriginal and Torres Strait Islander Social Survey (NATSISS) or National Aboriginal and Torres Strait Islander Health Survey (NATSIHS) data may be used for analysis, dependent upon which survey is most recent.

Data on persons 'at risk of long term alcohol related harm' had previously been defined according to the 2001 NHMRC guidelines: for males, an average of more than 4 standard drinks per day in the last week, and for females, an average of more than 2 standard drinks per day in the last week.

Representational attributes

Representation class: Percentage

Data type: Real
Unit of measure: Person

Format: N[N].N

Indicator conceptual framework

Framework and dimensions:

Health behaviours

Data source attributes

Data sources:

Data Source

ABS 2014-15 National Health Survey (NHS)

Frequency

Every 3 years

Data custodian

Australian Bureau of Statistics

Data Source

ABS 2014-15 National Aboriginal and Torres Strait Islander Social Survey (NATSISS)

Frequency

Every 6 years

Data custodian

Australian Bureau of Statistics

Accountability attributes

Reporting requirements: National Healthcare Agreement

Organisation responsible

for providing data:

Australian Bureau of Statistics

Further data development / Specification: Final, the measure meets the intention of the indicator.

collection required:

Relational attributes

Related metadata references:

Supersedes <u>National Healthcare Agreement: PI 05–Levels of risky alcohol consumption, 2016</u>

Health!, Superseded 31/01/2017

Has been superseded by <u>National Healthcare Agreement: PI 05–Levels of risky</u> <u>alcohol consumption, 2018</u>

Health!, Superseded 19/06/2019

See also National Healthcare Agreement: PB e—Better health: by 2018, reduce the national smoking rate to 10 per cent of the population and halve the Indigenous smoking rate over the 2009 baseline, 2017

Health!, Superseded 30/01/2018

See also National Healthcare Agreement: PI 03—Prevalence of overweight and obesity, 2017

Health!, Superseded 30/01/2018

See also National Healthcare Agreement: PI 04–Rates of current daily smokers, 2017

Health!, Superseded 30/01/2018

See also National Healthcare Agreement: PI 16—Potentially avoidable deaths, 2017

Health!, Superseded 30/01/2018

See also National Indigenous Reform Agreement: PI 04-Levels of risky alcohol consumption, 2018

Indigenous, Superseded 31/07/2018