

Australian Atlas of Healthcare Variation: Number of potentially preventable hospitalisations - kidney and urinary tract infections per 100,000 people, 2014-15

Exported from METEOR (AIHW's Metadata Online Registry)

© Australian Institute of Health and Welfare 2024

This product, excluding the AIHW logo, Commonwealth Coat of Arms and any material owned by a third party or protected by a trademark, has been released under a Creative Commons BY 4.0 (CC BY 4.0) licence. Excluded material owned by third parties may include, for example, design and layout, images obtained under licence from third parties and signatures. We have made all reasonable efforts to identify and label material owned by third parties.

You may distribute, remix and build on this website's material but must attribute the AIHW as the copyright holder, in line with our attribution policy. The full terms and conditions of this licence are available at <https://creativecommons.org/licenses/by/4.0/>.

Enquiries relating to copyright should be addressed to info@aihw.gov.au.

Enquiries or comments on the METEOR metadata or download should be directed to the METEOR team at meteor@aihw.gov.au.

Australian Atlas of Healthcare Variation: Number of potentially preventable hospitalisations - kidney and urinary tract infections per 100,000 people, 2014-15

Identifying and definitional attributes

| | |
|-----------------------------|--|
| Metadata item type: | Indicator |
| Indicator type: | Indicator |
| Short name: | Kidney and urinary tract infection hospitalisations, 2014-15 |
| METEOR identifier: | 638929 |
| Registration status: | Australian Commission on Safety and Quality in Health Care, Standard 07/06/2017 |
| Description: | Number of potentially preventable hospitalisations for kidney and urinary tract infections per 100,000 people, age and sex standardised. Data are disaggregated by the area in which the person lives. |
| Indicator set: | Australian Atlas of Healthcare Variation 2017 Australian Commission on Safety and Quality in Health Care, Standard 07/06/2017 |

Collection and usage attributes

| | |
|-----------------------------------|---|
| Population group age from: | All ages |
| Computation description: | Inclusion codes, description and additional requirements |

| ICD-10-AM 8th edition code | ICD-10-AM Description | Additional requirements |
|----------------------------|--|--------------------------|
| N10 | Acute tubulo-interstitial nephritis | Principal diagnosis only |
| N11.0 | Nonobstructive reflux-associated pyelonephritis | |
| N11.1 | Chronic obstructive pyelonephritis | |
| N11.8 | Other chronic tubulo-interstitial nephritis | |
| N11.9 | Chronic tubule-interstitial nephritis, unspecified | |
| N12 | Tubulo-interstitial nephritis, not specified as acute or chronic | |
| N13.6 | Pyonephrosis | |
| N15.1 | Renal and perinephric abscess | |
| N15.9 | Renal tubulo-interstitial disease, unspecified | |
| N28.9 | Disorders of kidney and ureter, unspecified | |
| N39.0 | Urinary tract infection, site not specified | |
| N39.9 | Disorder of urinary system, unspecified | |

Exclusion codes and description

| Care type | Description |
|-----------|-------------------------------|
| 7.3 | Newborn—unqualified days only |
| 9.0 | Organ procurement—posthumous |
| 10.0 | Hospital boarder |

Presented as a number per 100,000 people.

Rates for all Australians are directly age- and sex- standardised, to the 2001 Australian population, using 5-year age groups: 0-4, 5-9, ... , 80-84, 85 and over.

Indigenous and other Australian rates are directly age and sex standardised, to the 2001 Australian population, using 5-year age groups: 0-4, 5-9, ... , 60-64, 65 and over.

For more information about age-standardisation in general see </content/index.phtml/itemId/327276>

Analysis by Statistical Area Level 3 (SA3) is based on Statistical Area Level 2 (SA2) of usual residence of the patient.

Suppress data (number or rate) if at least one of the following conditions are met:

- the total denominator is less than 1,000
- the total numerator is less than 20

Age and sex standardised rates are suppressed where the denominator for at least one of the age and sex groups used to calculate the rate is below 30.

Computation:

$100,000 \times (\text{Numerator} \div \text{Denominator})$

Numerator:

Number of kidney and urinary tract infection hospitalisations, 2014-15.

Numerator data elements:

Data Element / Data Set

[Person—date of birth, DDMMYYYY](#)

NMDS / DSS

[Admitted patient care NMDS 2014-15](#)

Guide for use

Data source type: Administrative by-product data

Data Element / Data Set

[Hospital service—care type, code N\[N\]](#)

NMDS / DSS

[Admitted patient care NMDS 2014-15](#)

Guide for use

Data source type: Administrative by-product data

Data Element / Data Set

[Episode of care—principal diagnosis, code \(ICD-10-AM 8th edn\) ANN{,N\[N\]}](#)

Data Source

[National Hospital Morbidity Database \(NHMD\)](#)

NMDS / DSS

[Admitted patient care NMDS 2014-15](#)

Denominator:

Total population as at 30 June, 2014.

Denominator data elements:

Data Element / Data Set

Person—estimated resident population of Australia

Data Source

[ABS Indigenous experimental estimates and projections \(2001 Census-based\)](#)

Guide for use

Data source type: Census based plus administrative by-product data

Data Element / Data Set

[Person—estimated resident population of Australia, total people N\[N\(7\)\]](#)

Data Source

[ABS Estimated resident population \(total population\)](#)

Guide for use

Disaggregation:

SA3 by:

- remoteness (ASGC Remoteness structure) and Socio-Economic Indexes for Areas (SEIFA) Index of Relative Socioeconomic Disadvantage (IRSD)

State and territory by Indigenous status

Disaggregation data elements:**Data Element / Data Set**

[Person—area of usual residence, statistical area level 2 \(SA2\) code \(ASGS 2011\) N\(9\)](#)

Data Source

[National Hospital Morbidity Database \(NHMD\)](#)

Representational attributes

Representation class: Rate
Data type: Integer
Unit of measure: Episode
Format: N[NNNN]

Data source attributes

Data sources:**Data Source**

[ABS Indigenous experimental estimates and projections \(2001 Census-based\)](#)

Frequency

Periodic

Data quality statement

[ABS Indigenous experimental estimates and projections, QS](#)

Data custodian

Australian Bureau of Statistics

Data Source

[National Hospital Morbidity Database \(NHMD\)](#)

Frequency

Annual

Data custodian

Australian Institute of Health and Welfare

Data Source

[ABS Estimated resident population \(total population\)](#)

Frequency

Quarterly

Data quality statement

[ABS Estimated resident population \(total population\), QS](#)

Data custodian

Australian Bureau of Statistics

Accountability attributes**Methodology:**

Statistical Area Level 3 (SA3s) are geographic areas defined in the ABS Australian Statistical Geography Standard (ASGS). The aim of SA3s is to create a standard framework for the analysis of ABS data at the regional level through clustering groups of SA2s that have similar regional characteristics. There are 333 spatial SA3s covering the whole of Australia without gaps or overlaps. They are designed to provide a regional breakdown of Australia. SA3s generally have a population of between 30,000 and 130,000 people. There are approximately 50 with fewer than 30,000 people and 35 with more than 130,000 as at 30 June 2011. For further information see the ABS publication, Australian Statistical Geography Standard (ASGS): Volume 1 – Main Structure and Greater Capital City Statistical Areas, July 2011 (cat. no. 1270.0.55.001).

The scope of the NHMD is episodes of care for admitted patients in all public and private acute and psychiatric hospitals, free-standing day hospital facilities and alcohol and drug treatment centres in Australia. Hospitals operated by the Australian Defence Force, corrections authorities and in Australia's off-shore territories are not in scope, but some are included.

Private hospitals include private free-standing day hospital facilities and other private hospitals (which also include private psychiatric hospitals).

Reporting requirements: Australian Commission on Safety and Quality in Health Care
Australian Atlas of Healthcare Variation

Organisation responsible for providing data: Australian Institute of Health and Welfare

Accountability: Australian Commission on Safety and Quality in Health Care

Release date: 07/06/2017

Source and reference attributes

Submitting organisation: Australian Commission on Safety and Quality in Health Care

Reference documents: For more information about potentially preventable hospitalisations see:
National Healthcare Agreement: PI 18–Selected potentially preventable hospitalisations, 2017
[/content/index.phtml/itemId/630028](#)
[Australian Commission on Safety and Quality in Health Care and Australian Institute of Health and Welfare. Exploring Healthcare Variation in Australia: Analyses Resulting from an OECD Study. Sydney: ACSQHC, 2014.](#)
National Health Performance Authority 2015, Healthy Communities: Potentially preventable hospitalisations in 2013–14, Technical Supplement.

Relational attributes

Related metadata references: See also [Number of potentially preventable hospitalisations - kidney and urinary tract infections per 100,000 people of all ages, 2014-15 to 2017-18](#)
[Australian Commission on Safety and Quality in Health Care, Standard 27/04/2021](#)