# National Indigenous Reform Agreement: PI 07-Proportion of babies born of low birthweight, 2018

Proportion of babies born of low birthweight, 2018
Exported from METEOR (AIHW's Metadata Online Registry)

© Australian Institute of Health and Welfare 2024

This product, excluding the AlHW logo, Commonwealth Coat of Arms and any material owned by a third party or protected by a trademark, has been released under a Creative Commons BY 4.0 (CC BY 4.0) licence. Excluded material owned by third parties may include, for example, design and layout, images obtained under licence from third parties and signatures. We have made all reasonable efforts to identify and label material owned by third parties.

You may distribute, remix and build on this website's material but must attribute the AlHW as the copyright holder, in line with our attribution policy. The full terms and conditions of this licence are available at https://creativecommons.org/licenses/by/4.0/.

Enquiries relating to copyright should be addressed to info@aihw.gov.au.

Enquiries or comments on the METEOR metadata or download should be directed to the METEOR team at meteor@aihw.gov.au.

# National Indigenous Reform Agreement: PI 07-Proportion of babies born of low birthweight, 2018

### Identifying and definitional attributes

Metadata item type: Indicator Indicator type: Indicator

**Short name:** PI 07-Proportion of babies born of low birthweight, 2018

METEOR identifier: 668674

**Registration status:** <u>Indigenous</u>, Superseded 31/07/2018

**Description:** The incidence of low birthweight among live born singleton babies by Indigenous

status of mothers, and among live born singleton babies by Indigenous status of

babies.

Rationale: Low birthweight is associated with increased risk of poor health and death during

infancy and increased prevalence of a number of chronic diseases in adulthood.

Low birthweight is a particular issue for Indigenous Australians.

Indicator set: National Indigenous Reform Agreement (2018)

Indigenous, Superseded 31/07/2018

Outcome area: <a href="Indigenous children">Indigenous children are born and remain healthy</a>

Indigenous, Standard 21/07/2010

Data quality statement: National Indigenous Reform Agreement: PI 07-Proportion of babies born of low

birthweight, 2018; Quality Statement Indigenous, Superseded 07/02/2019

### Collection and usage attributes

#### Computation description:

Rates are calculated for Indigenous and non-Indigenous Australians.

Rate ratios and rate differences are calculated for comparisons between Indigenous and non-Indigenous Australians.

#### For variability bands:

Variability bands are to be calculated for rates (single year data and for total data for 3 years combined) using the standard method (see definition below).

#### For trends:

Percentage change and statistical significance of change is to be calculated (required for reporting of progress over time).

#### Presentation:

Number, percentage, rate per 100 babies, rate ratio, rate difference and variability bands.

Low birthweight is defined as less than 2,500 grams.

Excludes multiple births, stillbirths and births of less than 20 weeks gestation (where gestation is known).

Analysis by state/territory and remoteness is based on the usual residence of the mother.

Data exclude Australian non-residents, residents of external territories and records where state/territory of residence was not stated.

#### **Definitions:**

Standard method for variability band computation:

Rates derived from administrative data counts are not subject to sampling error but may still be subject to natural random variation, especially for small counts. A 95% confidence interval (CI) for an estimate is a range of values which is very likely (95 times out of 100) to contain the true unknown value. Where the 95% CIs of two estimates do not overlap it can be concluded that there is a statistically significant difference between the two estimates. This is the standard method used in Australian Institute of Health and Welfare (AIHW) publications for which formulas can be sourced from Breslow and Day (1987) in the publication *Statistical methods in cancer research*. Typically in the standard method, the observed rate is assumed to have natural variability in the numerator count (for example, deaths, hospital visits) but not in the population denominator count. Also, the rate is assumed to have been generated from a normal distribution ('Bell curve'). Random variation in the numerator count is assumed to be centred around the true value; that is, there is no systematic bias.

**Computation:** Rate: 100 x (Numerator ÷ Denominator).

Rate ratio: Indigenous rate divided by non-Indigenous rate.

Rate difference: Indigenous rate minus non-Indigenous rate.

**Variability bands:** to be calculated using the standard method for estimating 95% Cls as follows:

Crude rate:

$$CI(CR)_{95\%} = CR \pm 100 \times 1.96 \times \sqrt{\frac{\frac{CR}{100} \left(1 - \frac{CR}{100}\right)}{n}}$$

Where CI = confidence interval

CR = crude rate (expressed as a percentage)

n=number of live born singleton babies.

**Percentage change:** Calculated by multiplying the average annual change over the period by the number of data points less 1. This is then divided by the rate for the first year in the series and multiplied by 100.

The average annual change in rates, rate ratios and rate differences are calculated using linear regression which uses the least squares method to calculate a straight line that best fits the data and returns an array that best describes the line. The simple linear regression line, Y = a + bX, 'slope' (b) estimate was used to determine the average annual change in the data over the period. The formula used to calculate the slope estimate and standard error of the slope in Microsoft Excel is:

LINEST: (known\_y's, known\_x's, true) entered as an array formula (Ctrl, Shift, Enter).

**Statistical significance of change:** The 95% CIs for the standard error of the slope estimate (average annual change) are used to determine whether the apparent increases or decreases in the data are statistically significant at the p<0.05 level. The formula used to calculate the CIs for the standard error of the slope estimate is:

$$95\% \text{ CI}(x) = x \pm 1.96 \text{ x SE}(x)$$

where x is the average annual change (slope estimate). If the upper and lower 95% Cls do not include zero, then it can be concluded that there is statistical evidence of an increasing or decreasing trend in the data over the study period.

**Numerator:** Number of low birthweight live born singleton babies.

#### Numerator data elements:

-Data Element / Data Set-

Birth—birth weight, total grams NNNN

**Data Source** 

AlHW National Perinatal Data Collection (NPDC)

NMDS / DSS

Perinatal NMDS 2014-18

Guide for use

Data source type: Administrative by-product data

#### Data Element / Data Set-

Birth—birth status, code N

**Data Source** 

AlHW National Perinatal Data Collection (NPDC)

NMDS / DSS

Perinatal NMDS 2014-18

Guide for use

Data source type: Administrative by-product data

#### Data Element / Data Set-

Birth event—birth plurality, code N

**Data Source** 

AlHW National Perinatal Data Collection (NPDC)

NMDS / DSS

Perinatal NMDS 2014-18

Guide for use

Data source type: Administrative by-product data.

**Denominator:** 

Number of live-born singleton infants.

# Denominator data elements:

#### Data Element / Data Set-

Birth—birth status, code N

**Data Source** 

AlHW National Perinatal Data Collection (NPDC)

NMDS / DSS

Perinatal NMDS 2014-18

Guide for use

Data source type: Administrative by-product data.

#### Data Element / Data Set-

Birth event—birth plurality, code N

**Data Source** 

AlHW National Perinatal Data Collection (NPDC)

NMDS / DSS

Perinatal NMDS 2014-18

Guide for use

Data source type: Administrative by-product data.

#### Data Element / Data Set-

Product of conception—gestational age, completed weeks N[N]

**Data Source** 

AlHW National Perinatal Data Collection (NPDC)

NMDS / DSS

Perinatal NMDS 2014-18

Guide for use

Data source type: Administrative by-product data

#### Disaggregation:

#### Indigenous status of the mother

Current period—(2013-2015):

For Indigenous and non-Indigenous (number, rate, rate ratio, rate difference and variability bands):

- Total and state/territory: by Indigenous status of the mother.
- Total by remoteness area: by Indigenous status of the mother.

Time series—2007, 2008, 2009, 2010, 2011, 2012, 2013, 2014 (the data for these years have been previously supplied), 2015 (required for 2018 reporting):

For Indigenous and non-Indigenous (number, rate, rate ratio, rate difference and variability bands):

- Total and state/territory: by Indigenous status of the mother.
- Total by remoteness area (from 2012 onwards): by Indigenous status of the mother.

#### Indigenous status of the baby

Current period—(2014–2015):

For Indigenous and non-Indigenous (number, rate, rate ratio, rate difference and variability bands):

- Total and state/territory: by Indigenous status of the baby.
- Total by remoteness area: by Indigenous status of the baby.

Time series—2013, 2014 (these data have been previously supplied), 2015 (required for 2018 reporting).

For Indigenous and non-Indigenous (number, rate, rate ratio, rate difference and variability bands):

- Total and state/territory: by Indigenous status of the baby.
- Total by remoteness area: by Indigenous status of the baby.

# Indigenous status of the baby cross-tabulated by the Indigenous status of the mother

Current period only—(2014-2015):

For Indigenous and non-Indigenous (number, rate, rate ratio and rate difference):

• Total and state/territory: by Indigenous status of the baby and of the mother.

# Disaggregation data elements:

#### Data Element / Data Set-

Person-Indigenous status, code N

**Data Source** 

AlHW National Perinatal Data Collection (NPDC)

NMDS / DSS

Perinatal NMDS 2014-18

Guide for use

Data source type: Administrative by-product data.

#### Data Element / Data Set-

Person—area of usual residence, statistical area level 2 (SA2) code (ASGS 2011) N(9)

**Data Source** 

AlHW National Perinatal Data Collection (NPDC)

NMDS / DSS

Perinatal NMDS 2014-18

Guide for use

Data source type: Administrative by-product data

Used for disaggregation by state/territory and remoteness area. Classifications for remoteness area are based on Australian Standard Geographical Classification (ASGC) prior to 2012 and Australian Statistical Geography Standard (ASGS) from 2012.

#### Comments:

Most recent data available for the 2018 National Indigenous Reform Agreement (NIRA) Report (2016–17 reporting cycle) is 2015.

Aggregated data for a 3-year period (2013 to 2015) will be reported for the current reporting period.

Single year data (2007, 2008, 2009, 2010, 2011, 2012, 2013, 2014 and 2015) will be reported for time series, noting that previously supplied data will be used unless a resupply is provided.

Information is included in the National Perinatal Data Collection (NPDC) for all live births and stillbirths of at least 400 grams birthweight or at least 20 weeks gestation.

Until 2011, the NPDC only included information on the Indigenous status of the mother. From 2011, the NPDC included information on the Indigenous status of the baby for selected jurisdictions—New South Wales, Victoria, Queensland, the Australian Capital Territory and the Northern Territory. The mandatory collection of this data item began from July 2012 following its inclusion in the Perinatal National Minimum Data Set (NMDS). National data based on the NMDS data item are therefore available for the latter 6 months of 2012, and 2013 is the first full calendar year for which nationally standardised data on the Indigenous status of babies are available.

Disaggregation by Indigenous status is recommended to be reported using 3-year combined data for the current reporting period due to the relatively small number of low birthweight babies born to Indigenous mothers each year. Single year data are to be reported for time series.

To report trends, the body assessing progress over time may separately request percentage change and statistical significance testing for this indicator directly from the data provider (AlHW).

Variability bands accompanying perinatal data should be used where there are at least 3 full calendar years of data available for the purposes of comparisons over time and for national estimates at a point in time for Indigenous/non-Indigenous comparisons. Variability bands will be provided for Indigenous status of the baby stratified by Indigenous status of the mother for the first time, as this is the first year of reporting 3 full years of data.

Baseline year for the Council of Australian Governments' NIRA (Halve the gap in mortality rates for Indigenous children under 5 within a decade) is 2008; baseline year for this indicator is 2007; target year is 2018.

The term 'Aboriginal and Torres Strait Islander people' is preferred when referring to the separate Indigenous peoples of Australia. However, the term 'Indigenous' is used interchangeably with 'Aboriginal and Torres Strait Islander' in this indicator set to assist readability.

## Representational attributes

Representation class: Percentage

Data type:RealUnit of measure:PersonFormat:N[N].N

## Indicator conceptual framework

Framework and dimensions:

**Health Conditions** 

Data source attributes

Data sources: **Data Source** 

AlHW National Perinatal Data Collection (NPDC)

Frequency

Calendar years ending 31 December each year

Data custodian

Australian Institute of Health and Welfare

### Accountability attributes

Reporting requirements: National Indigenous Reform Agreement.

Organisation responsible

for providing data:

Australian Institute of Health and Welfare

collection required:

Further data development / A data item for Indigenous status of the baby was added to the Perinatal National Minimum Data Set (NMDS) for collection from July 2012 onwards.

### Source and reference attributes

**Submitting organisation:** Australian Institute of Health and Welfare

Steward: National Indigenous Reform Agreement Performance Information Management

Group

Reference documents: Breslow NE & Day NE (eds) 1987. Statistical methods in cancer research. Volume

II: The design and analysis of cohort studies. IARC Scientific Publications No. 82. Lyon, France: International Agency for Research on Cancer. Viewed 20 June 2017,

http://www.iarc.fr/en/publications/pdfs-online/stat/sp82/

### Relational attributes

Related metadata references:

Supersedes National Indigenous Reform Agreement: PI 07—Proportion of babies

born of low birthweight, 2017

Indigenous, Superseded 06/06/2017

Has been superseded by National Indigenous Reform Agreement: PI 07-Proportion

of babies born of low birthweight, 2019 Indigenous, Superseded 23/08/2019

See also National Healthcare Agreement: PI 01-Proportion of babies born of low

birth weight, 2017

Health!, Superseded 30/01/2018