

Episode of care—principal diagnosis, code (ICD-10-AM 10th edn) ANN{.N[N]}

Exported from METEOR (AIHW's Metadata Online Registry)

© Australian Institute of Health and Welfare 2024

This product, excluding the AIHW logo, Commonwealth Coat of Arms and any material owned by a third party or protected by a trademark, has been released under a Creative Commons BY 4.0 (CC BY 4.0) licence. Excluded material owned by third parties may include, for example, design and layout, images obtained under licence from third parties and signatures. We have made all reasonable efforts to identify and label material owned by third parties.

You may distribute, remix and build on this website's material but must attribute the AIHW as the copyright holder, in line with our attribution policy. The full terms and conditions of this licence are available at <https://creativecommons.org/licenses/by/4.0/>.

Enquiries relating to copyright should be addressed to info@aihw.gov.au.

Enquiries or comments on the METEOR metadata or download should be directed to the METEOR team at meteor@aihw.gov.au.

Episode of care—principal diagnosis, code (ICD-10-AM 10th edn) ANN{.N[N]}

Identifying and definitional attributes

Metadata item type:	Data Element
Short name:	Principal diagnosis
METEOR identifier:	680976
Registration status:	Health! , Superseded 12/12/2018 Tasmanian Health , Superseded 08/04/2019
Definition:	The diagnosis established after study to be chiefly responsible for occasioning an episode of admitted patient care, an episode of residential care or an attendance at the health care establishment, as represented by a code.
Data Element Concept:	Episode of care—principal diagnosis
Value Domain:	Diagnosis code (ICD-10-AM 10th edn) ANN{.N[N]}

Value domain attributes

Representational attributes

Classification scheme:	International Statistical Classification of Diseases and Related Health Problems, Tenth Revision, Australian Modification 10th edition
Representation class:	Code
Data type:	String
Format:	ANN{.N[N]}
Maximum character length:	6

Data element attributes

Collection and usage attributes

Guide for use:	<p>The principal diagnosis must be determined in accordance with the Australian Coding Standards. Each episode of admitted patient care must have a principal diagnosis and may have additional diagnoses. The diagnosis can include a disease, condition, injury, poisoning, sign, symptom, abnormal finding, complaint, or other factor influencing health status.</p> <p>As a minimum requirement the Principal diagnosis code must be a valid code from the current edition of the <i>International statistical classification of diseases and related health problems, 10th revision, Australian modification</i> (ICD-10-AM).</p> <p>For episodes of admitted patient care, some diagnosis codes are too imprecise or inappropriate to be acceptable as a principal diagnosis and will group to an error DRG in the Australian Refined Diagnosis Related Groups.</p> <p>Diagnosis codes starting with a V, W, X or Y, describing the circumstances that cause an injury, rather than the nature of the injury, cannot be used as a principal diagnosis. Diagnosis codes which are morphology codes cannot be used as a principal diagnosis.</p>
Collection methods:	A principal diagnosis should be recorded and coded upon separation , for each episode of admitted patient care or episode of residential care or attendance at a health-care establishment. The principal diagnosis is derived from and must be substantiated by clinical documentation.

Comments: The principal diagnosis is one of the most valuable health data elements. It is used for epidemiological research, casemix studies and planning purposes.

Source and reference attributes

Origin: Australian Consortium for Classification Development
National Data Standard for Injury Surveillance Advisory Group

Relational attributes

Related metadata references: Supersedes [Episode of care—principal diagnosis, code \(ICD-10-AM 10th edn\) ANN{,N\[N\]}](#)
[Health!](#), Superseded 25/01/2018
[Independent Hospital Pricing Authority](#), Recorded 04/08/2016
[Tasmanian Health](#), Superseded 08/04/2019

Has been superseded by [Episode of care—principal diagnosis, code \(ICD-10-AM 11th edn\) ANN{,N\[N\]}](#)
[Health!](#), Superseded 20/10/2021
[Tasmanian Health](#), Standard 08/04/2019

Is used in the formation of [Episode of admitted patient care—diagnosis related group, code \(AR-DRG v 9.0\) ANNA](#)
[Tasmanian Health](#), Superseded 19/06/2020

See also [Episode of care—additional diagnosis, code \(ICD-10-AM 10th edn\) ANN{,N\[N\]}](#)
[Health!](#), Superseded 12/12/2018
[Tasmanian Health](#), Superseded 08/04/2019

Implementation in Data Set Specifications: [Activity based funding: Mental health care NBEDS 2018-19](#)
[Health!](#), Superseded 12/12/2018
Implementation start date: 01/07/2018
Implementation end date: 30/06/2019

[Admitted patient care NMDS 2018-19](#)
[Health!](#), Superseded 12/12/2018
Implementation start date: 01/07/2018
Implementation end date: 30/06/2019
Conditional obligation:

The principal diagnosis is a major determinant in the classification of Australian Refined Diagnosis Related Groups and Major Diagnostic Categories.

Where the principal diagnosis is recorded prior to discharge (as in the annual census of public psychiatric hospital patients), it is the current provisional principal diagnosis. Only use the admission diagnosis when no other diagnostic information is available. The current provisional diagnosis may be the same as the admission diagnosis.

[Community mental health care NMDS 2018-19](#)
[Health!](#), Superseded 12/12/2018
Implementation start date: 01/07/2018
Implementation end date: 30/06/2019

[Residential mental health care NMDS 2018-19](#)
[Health!](#), Superseded 12/12/2018
Implementation start date: 01/07/2018
Implementation end date: 30/06/2019
DSS specific information:

Codes can be used from ICD-10-AM or from The ICD-10-AM Mental Health Manual: An Integrated Classification and Diagnostic Tool for Community-Based Mental Health Services, published by the National Centre for Classification in Health 2002.

The principal diagnosis should be recorded and coded upon the end of an episode of residential care (i.e. annually for continuing residential care).

Implementation in Indicators:

Used as Numerator

[Australian Health Performance Framework: PI 2.1.4–Selected potentially preventable hospitalisations, 2020](#)

[Health!](#), Standard 01/12/2020

[Australian Health Performance Framework: PI 3.1.1–Incidence of heart attacks \(acute coronary events\), 2020](#)

[Health!](#), Standard 13/10/2021

[Australian Health Performance Framework: PI 3.1.5–Hospitalisation for injury and poisoning, 2020](#)

[Health!](#), Standard 13/10/2021

[Heavy menstrual bleeding clinical care standard indicators: 8-Hospital rate of hysterectomy per 100 episodes](#)

[Health!](#), Standard 17/10/2018

[National Healthcare Agreement: PB f–By 2014–15, improve the provision of primary care and reduce the proportion of potentially preventable hospital admissions by 7.6 per cent over the 2006–07 baseline to 8.5 per cent of total hospital admissions, 2021](#)

[Health!](#), Standard 16/09/2020

[National Healthcare Agreement: PI 09–Incidence of heart attacks \(acute coronary events\), 2021](#)

[Health!](#), Standard 16/09/2020

[National Healthcare Agreement: PI 09–Incidence of heart attacks \(acute coronary events\), 2022](#)

[Health!](#), Standard 24/09/2021

[National Healthcare Agreement: PI 18–Selected potentially preventable hospitalisations, 2021](#)

[Health!](#), Standard 16/09/2020

[National Healthcare Agreement: PI 23–Unplanned hospital readmission rates, 2021](#)

[Health!](#), Standard 19/11/2020

[National Healthcare Agreement: PI 27–Number of hospital patient days used by those eligible and waiting for residential aged care, 2021](#)

[Health!](#), Standard 16/09/2020

[Number of lumbar spinal decompression \(excluding lumbar spinal fusion\) hospitalisations per 100,000 people aged 18 years and over, 2012-13 to 2014-15 and 2015-16 to 2017-18](#)

[Australian Commission on Safety and Quality in Health Care](#), Standard 27/04/2021

[Number of lumbar spinal fusion \(excluding lumbar spinal decompression\) hospitalisations per 100,000 people, aged 18 years and over, 2012-13 to 2014-15 and 2015-16 to 2017-18](#)

[Australian Commission on Safety and Quality in Health Care](#), Standard 27/04/2021

[Number of lumbar spinal fusion \(with or without lumbar spinal decompression\) hospitalisations per 100,000 people, aged 18 years and over, 2012-13 to 2014-15 and 2015-16 to 2017-18](#)

[Australian Commission on Safety and Quality in Health Care](#), Standard 27/04/2021

Used as Disaggregation

[Number of lumbar spinal decompression \(excluding lumbar spinal fusion\) hospitalisations per 100,000 people aged 18 years and over, 2012-13 to 2014-15 and 2015-16 to 2017-18](#)

[Australian Commission on Safety and Quality in Health Care](#), Standard 27/04/2021

Used as Denominator

[Heavy menstrual bleeding clinical care standard indicators: 8-Hospital rate of hysterectomy per 100 episodes](#)

[Health!](#), Standard 17/10/2018

[Third and Fourth Degree Perineal Tears Clinical Care Standard: 3b-Proportion of women who had an instrumental vaginal birth using forceps](#)

