

# Female—postpartum perineal status, code N[N]

Exported from METEOR (AIHW's Metadata Online Registry)

© Australian Institute of Health and Welfare 2024

This product, excluding the AIHW logo, Commonwealth Coat of Arms and any material owned by a third party or protected by a trademark, has been released under a Creative Commons BY 4.0 (CC BY 4.0) licence. Excluded material owned by third parties may include, for example, design and layout, images obtained under licence from third parties and signatures. We have made all reasonable efforts to identify and label material owned by third parties.

You may distribute, remix and build on this website's material but must attribute the AIHW as the copyright holder, in line with our attribution policy. The full terms and conditions of this licence are available at <https://creativecommons.org/licenses/by/4.0/>.

Enquiries relating to copyright should be addressed to [info@aihw.gov.au](mailto:info@aihw.gov.au).

Enquiries or comments on the METEOR metadata or download should be directed to the METEOR team at [meteor@aihw.gov.au](mailto:meteor@aihw.gov.au).

# Female—postpartum perineal status, code N[N]

## Identifying and definitional attributes

<b>Metadata item type:</b>	Data Element
<b>Short name:</b>	Postpartum perineal status
<b>METEOR identifier:</b>	717924
<b>Registration status:</b>	<a href="#">Health!</a> , Superseded 03/12/2020
<b>Definition:</b>	The state of a female's perineum following birth, as represented by a code.
<b>Context:</b>	Perinatal
<b>Data Element Concept:</b>	<a href="#">Female—postpartum perineal status</a>
<b>Value Domain:</b>	<a href="#">Perineal status code N[N]</a>

## Value domain attributes

### Representational attributes

<b>Representation class:</b>	Code
<b>Data type:</b>	Number
<b>Format:</b>	N[N]
<b>Maximum character length:</b>	2

	<b>Value</b>	<b>Meaning</b>
<b>Permissible values:</b>	1	Intact
	2	1st degree laceration/vaginal graze
	3	2nd degree laceration
	4	3rd degree laceration
	5	Episiotomy
	7	4th degree laceration
	88	Other perineal laceration, rupture or tear
<b>Supplementary values:</b>	99	Not stated/inadequately described

## Collection and usage attributes

**Guide for use:****CODE 1 Intact**

Should not be recorded in conjunction with any other code.

**CODE 2 1st degree laceration/vaginal graze**

Graze, laceration, rupture or tear of the perineal skin during delivery that may be considered to be slight or that involves one or more of the following structures:

- fourchette
- labia
- periurethral tissue (excluding involvement of urethra)
- vagina, low
- skin
- vulva.

Should not be recorded in conjunction with any other code except for Code 5 (Episiotomy).

**CODE 3 2nd degree laceration**

Perineal laceration, rupture or tear as in Code 2 occurring during delivery, also involving:

- pelvic floor
- perineal muscles
- vaginal muscles.

Excludes laceration involving the anal sphincter. Should not be recorded in conjunction with any other code except for Code 5 (Episiotomy).

**CODE 4 3rd degree laceration**

Perineal laceration, rupture or tear as in Code 3 occurring during delivery, also involving:

- anal sphincter (excluding involvement of anal or rectal mucosa)
- rectovaginal septum
- sphincter not otherwise specified (NOS).

Excludes laceration involving the anal or rectal mucosa. Should not be recorded in conjunction with any other code except for Code 5 (Episiotomy).

**CODE 7 4th degree laceration**

Perineal laceration, rupture or tear as in Code 4 occurring during delivery, also involving:

- anal mucosa
- rectal mucosa.

Should not be recorded in conjunction with any other code except for Code 5 (Episiotomy).

**CODE 88 Other perineal laceration, rupture or tear**

May include haematoma or unspecified perineal tear. Should not be recorded in conjunction with any other code except for Code 5 (Episiotomy).

**Comments:**

If a laceration occurred during delivery and an episiotomy was performed, both the degree of laceration and the episiotomy should be recorded. If an episiotomy is performed, the perineum cannot be intact.

While 4th degree laceration is more severe than an episiotomy, this category has not been placed in order of clinical significance within the value domain.

Code 6 has been omitted as it is no longer in use. For information about its meaning in previous data elements, see superseded versions.

## Source and reference attributes

**Submitting organisation:** Australian Institute of Health and Welfare

**Origin:** ACCD (Australian Consortium for Classification Development) 2019. The International Statistical Classification of Diseases and Related Health Problems, 10th Revision, Australian Modification (ICD-10-AM), 11th edn. Tabular list and Alphabetic index. Darlinghurst, NSW: Independent Hospital Pricing Authority.

## Data element attributes

### Collection and usage attributes

**Guide for use:** More than one permissible value may be recorded.

**Comments:** Perineal laceration (tear) may cause significant maternal morbidity in the postnatal period. Episiotomy is an indicator of management during labour and, to some extent, of obstetric intervention.

### Relational attributes

**Related metadata references:** Supersedes [Female—postpartum perineal status, code N\[N\] Health!](#), Superseded 20/11/2019

Has been superseded by [Female—postpartum perineal status, code N\[N\] Health!](#), Superseded 17/12/2021

**Implementation in Data Set Specifications:** [Perinatal NMDS 2020–21 Health!](#), Superseded 03/12/2020

**Implementation start date:** 01/07/2020

**Implementation end date:** 30/06/2021

**DSS specific information:**

This is a multiple response data element and is therefore operationalised for data collection across 7 individual data items (one data item per permissible value, excluding Code 99 (Not stated/inadequately described)).