Delirium clinical care standard indicators: 5a-Proportion of patients with delirium who had a comprehensive assessment that includes relevant multidisciplinary consultation to investigate the cause(s) of delirium



© Australian Institute of Health and Welfare 2024

This product, excluding the AIHW logo, Commonwealth Coat of Arms and any material owned by a third party or protected by a trademark, has been released under a Creative Commons BY 4.0 (CC BY 4.0) licence. Excluded material owned by third parties may include, for example, design and layout, images obtained under licence from third parties and signatures. We have made all reasonable efforts to identify and label material owned by third parties.

You may distribute, remix and build on this website's material but must attribute the AlHW as the copyright holder, in line with our attribution policy. The full terms and conditions of this licence are available at https://creativecommons.org/licenses/by/4.0/.

Enquiries relating to copyright should be addressed to info@aihw.gov.au.

Enquiries or comments on the METEOR metadata or download should be directed to the METEOR team at meteor@aihw.gov.au.

Delirium clinical care standard indicators: 5a-Proportion of patients with delirium who had a comprehensive assessment that includes relevant multidisciplinary consultation to investigate the cause(s) of delirium

Identifying and definitional attributes

Metadata item type: Indicator Indicator type: Indicator

Short name: 5a- Proportion of patients with delirium who had a comprehensive assessment that

includes relevant multidisciplinary consultation to investigate the cause(s) of

delirium

METEOR identifier: 745818

Registration status: Australian Commission on Safety and Quality in Health Care, Standard 09/09/2021

Description: Proportion of patients with delirium who had a comprehensive assessment that

includes relevant multidisciplinary consultation to investigate the cause(s) of

delirium.

Indicator set: Clinical care standard indicators: delirium 2021

Australian Commission on Safety and Quality in Health Care, Standard

09/09/2021

Collection and usage attributes

Computation description:

For the numerator and denominator identify patients with delirium using the ICD codes specified in the <u>Hospital-Acquired Complications</u> (HACs) <u>List-Specifications</u>, worksheet 1, where complication group = 11. Delirium.

The numerator includes patients with delirium for whom a comprehensive assessment was undertaken, in consultation with the patient and their family or carer where appropriate, to investigate the cause(s) of their delirium.

To be included in the numerator, the comprehensive assessment must include multidisciplinary consultation, undertaken during the hospital stay, be documented in the patient's medical record, and include all of the following (ACSQHC, 2021):

- A medical and social history paying close attention to the patient's medication history and their pain management needs, and their hydration and nutritional status
- A physical examination, and
- Investigations (blood tests, urine tests, etc.), informed by the patient's medical history and physical examination.

'Multidisciplinary' means the assessment includes consultations with clinicians from more than one discipline. These consultations can be in person or via telehealth.

Presented as a percentage.

Computation: (Numerator ÷ Denominator) x 100

Numerator: The number of patients with delirium who had a comprehensive assessment that

includes relevant multidisciplinary consultation to investigate the cause(s) of

delirium.

Denominator: The number of patients with delirium.

Representational attributes

Representation class: Percentage

Data type:RealUnit of measure:PersonFormat:N[NN]

Source and reference attributes

Submitting organisation: Australian Commission on Safety and Quality in Health Care

Reference documents: Australian Commission on Safety and Quality in Health Care. Delirium Clinical

Care Standard. Sydney: ACSQHC; 2021.